Subject Index for *Hydrobiological Journal*, Volume 31, 1995

Abiotic environmental factors, **5**, 101–106
Abramis brama, **3**, 46–53
Abundance of turbot eggs, **6**, 76–86
Abundance variation, **3**, 87–89
Actinulae behavior, **6**, 84–92
Actinulae settlement, **6**, 84–92
Algae, **2**, 33–39; 59–66
Algae growth, **3**, 35–45
Ammoscalariidae, **4**, 65–69
Ampharetidae, **3**, 90–96
Anatomy of gonads, **3**, 9–14
Angara–Yenisey basin, **6**, 93–106
Anopla subclass, **6**, 22–34
Anti-fouling compounds, **6**, 66–75
Aquatic communities, **5**, 36–43
Aquatic ecosystems assessment, **4**, 103–109
Aquatic environment, **1**, 76–83; **5**, 1–9
Aquatic organisms, **2**, 1–5
Arabian Sea, **7**, 1–3
Araeolaimida, **2**, 94–108
Arcellidae, **7**, 19–32
Arctic Ocean, **4**, 34–50
Arietellidae, **7**, 99–103
Arrowworm species, **4**, 1–11
Artemia salina, **1**, 29–35
Astacidae, **3**, 87–89
Attachment mechanism, **7**, 68–77
Axonolaimidae, **2**, 94–108

Bacteria, **2**, 59–66
Bacteriology, **2**, 49–58
Bacterioplankton role, **5**, 66–79
Baltic Sea gulfs, **5**, 51–65
Behavioral response, **6**, 66–75
Benthic waters, **2**, 23–32
Bioassay, **1**, 76–83
Biological contractors, **5**, 44–50
Biological improvement, **2**, 49–58
Bivalves, **6**, 66–75
Bivalvia, **3**, 1–8; **7**, 8–18
Black Sea, **1**, 66–75; 92–101; **2**, 33–39; **3**, 9–14; **5**, 30–35; 36–43; **6**, 76–86
Black Sea inshore zone, **1**, 45–54
Bony fishes, **1**, 55–65
Bottom sediments, **5**, 51–65
Bregmacerotidae, **7**, 1–3
Bunodera luciopercae, **6**, 18–21; 107–110
Bunoderidae, **6**, 18–21; 107–110

Calycophorae, **7**, 33–50
Caspian Sea, **3**, 87–89
Cenoecomorphic structure, **2**, 13–22
Centrohelidian helizoans, **4**, 51–64
Centroheliozoa, **4**, 51–64; **6**, 45–60
Centropyxidae, **7**, 19–32
Cercariae, **4**, 25–33
Cesium-137 accumulation, **1**, 84–91
Chaetognatha, **4**, 1–11
Chemical technique, **2**, 6–12
Chernaya River estuary, **4**, 12–19
Chernobyl, **3**, 46–53
Chromium, **6**, 9–17
Chromium concentrations, **6**, 9–17
Chromium forms, **6**, 9–17
Chromosomal mutations, **1**, 55–65
Cladocera, **6**, 93–106
Cladocerans, **6**, 61–65
Clupeonella cultriventris cultriventris, **4**, 92–102
Cnidaria, **7**, 33–50
Colonization, **1**, 36–44
Colossendeis, **7**, 78–84
Commensal polychaete, **3**, 97–113
Computerized identification keys, **2**, 1–5
Consumption of primary production, **2**, 40–48
Copepoda, **7**, 99–103
Corambidae, **7**, 59–67
Crayfish, **3**, 87–89
Crustacea, **6**, 79–83; **7**, 68–77; 99–103
Cucumaria japonica, **6**, 35–44

Subject Index for *Hydrobiological Journal*, Volume 31, 1995

Abiotic environmental factors, **5**, 101–106
Abramis brama, **3**, 46–53
Abundance of turbot eggs, **6**, 76–86
Abundance variation, **3**, 87–89
Actinulae behavior, **6**, 84–92
Actinulae settlement, **6**, 84–92
Algae, **2**, 33–39; 59–66
Algae growth, **3**, 35–45
Ammoscalariidae, **4**, 65–69
Ampharetidae, **3**, 90–96
Anatomy of gonads, **3**, 9–14
Angara–Yenisey basin, **6**, 93–106
Anopla subclass, **6**, 22–34
Anti-fouling compounds, **6**, 66–75
Aquatic communities, **5**, 36–43
Aquatic ecosystems assessment, **4**, 103–109
Aquatic environment, **1**, 76–83; **5**, 1–9
Aquatic organisms, **2**, 1–5
Arabian Sea, **7**, 1–3
Araeolaimida, **2**, 94–108
Arcellidae, **7**, 19–32
Arctic Ocean, **4**, 34–50
Arietellidae, **7**, 99–103
Arrowworm species, **4**, 1–11
Artemia salina, **1**, 29–35
Astacidae, **3**, 87–89
Attachment mechanism, **7**, 68–77
Axonolaimidae, **2**, 94–108

Bacteria, **2**, 59–66
Bacteriology, **2**, 49–58
Bacterioplankton role, **5**, 66–79
Baltic Sea gulfs, **5**, 51–65
Behavioral response, **6**, 66–75
Benthic waters, **2**, 23–32
Bioassay, **1**, 76–83
Biological contractors, **5**, 44–50
Biological improvement, **2**, 49–58
Bivalves, **6**, 66–75
Bivalvia, **3**, 1–8; **7**, 8–18
Black Sea, **1**, 66–75; 92–101; **2**, 33–39; **3**, 9–14; **5**, 30–35; 36–43; **6**, 76–86
Black Sea inshore zone, **1**, 45–54
Bony fishes, **1**, 55–65
Bottom sediments, **5**, 51–65
Bregmacerotidae, **7**, 1–3
Bunodera luciopercae, **6**, 18–21; 107–110
Bunoderidae, **6**, 18–21; 107–110

Calycophorae, **7**, 33–50
Caspian Sea, **3**, 87–89
Cenoecomorphic structure, **2**, 13–22
Centrohelidian helizoans, **4**, 51–64
Centroheliozoa, **4**, 51–64; **6**, 45–60
Centropyxidae, **7**, 19–32
Cercariae, **4**, 25–33
Cesium-137 accumulation, **1**, 84–91
Chaetognatha, **4**, 1–11
Chemical technique, **2**, 6–12
Chernaya River estuary, **4**, 12–19
Chernobyl, **3**, 46–53
Chromium, **6**, 9–17
Chromium concentrations, **6**, 9–17
Chromium forms, **6**, 9–17
Chromosomal mutations, **1**, 55–65
Cladocera, **6**, 93–106
Cladocerans, **6**, 61–65
Clupeonella cultriventris cultriventris, **4**, 92–102
Cnidaria, **7**, 33–50
Colonization, **1**, 36–44
Colossendeis, **7**, 78–84
Commensal polychaete, **3**, 97–113
Computerized identification keys, **2**, 1–5
Consumption of primary production, **2**, 40–48
Copepoda, **7**, 99–103
Corambidae, **7**, 59–67
Crayfish, **3**, 87–89
Crustacea, **6**, 79–83; **7**, 68–77; 99–103
Cucumaria japonica, **6**, 35–44

Cumulative toxicity, **1**, 76–83
Cyclocalyx, **7**, 8–18

Danube delta lagoons, **1**, 11–28
Daphnia species, **6**, 93–106
Daphniidae, **6**, 93–106; **7**, 68–77
Daphniiformes, **7**, 68–77
Decapoda, **6**, 79–83
Dendrochirota, **6**, 35–44
Different-age morphology, **6**, 35–44
Distribution of turbot eggs, **6**, 76–86
Diversity, **6**, 45–60
Dnieper reservoirs, **1**, 1–10; 84–91; **5**, 15–29; **6**, 9–17
Dnieper River reservoirs, **5**, 107–110
Dnieper-Bug Lagoon, **6**, 9–17
Dniester River, **3**, 70–75

Eastern Europe, **6**, 45–60
Eastern Murman, **3**, 15–29
Ecosystem, **3**, 54–69
Effect of environmental pH, **3**, 35–45
Enoplida, **6**, 76–78
Environment pH, **3**, 35–45
Ethology, **7**, 85–98
Evolution, **1**, 55–65
Evolution of free-living Turbellaria, **2**, 71–89

Fecundity variation, **6**, 107–110
Feeding rate, **7**, 85–98
Fish culture ponds, **2**, 40–48
Fish muscle tissue, **2**, 6–12
Foraminifera, **4**, 65–69
Free-living marine nematodes, **2**, 94–108

Gastrolepidia clavigera, **3**, 97–113
Gastropoda, **7**, 51–58; 59–67
Genera related to *Sosane*, **3**, 90–96
Genome mutations, **1**, 55–65
Genus *Choanocystis*, **6**, 45–60
Genus *Dinophysis*, **2**, 33–39
Genus *Dreissena*, **5**, 107–110
Genus *Malacoceros*, **4**, 34–50
Genus *Parodontophora*, **2**, 94–108
Genus revision, **7**, 33–50
Grass carp, **6**, 1–8
Grass-carp organs, **6**, 1–8
Grass-carp tissues, **6**, 1–8
Gulf of Kandalaksha, **4**, 12–19; 51–64

Halalaimus dolgovi sp. n., **4**, 20–24
Halonymphidae, **3**, 1–8
Head skeletal structures, **4**, 1–11
Hemoglobin concentration, **5**, 80–90
Higher taxa, **6**, 22–34
Highly specific parasites, **6**, 61–65
Holothuriidae, **3**, 97–113
Holothurioidea, **3**, 97–113
Holothuroid, **6**, 35–44
Holothuroid hosts, **3**, 97–113
Holothuroidea, **6**, 35–44
Human impact, **1**, 11–28
Hyalospheniidae, **7**, 19–32
Hydrichthys, **7**, 1–3
Hydrida, **7**, 85–98
Hydrobiological indices, **4**, 103–109
Hydrocoryne, **2**, 67–70
Hydrocorynidae, **2**, 67–70
Hydroecological problems priority, **3**, 30–34
Hydroid polyps, **6**, 66–75

Inorganic nitrogen concentrations, **1**, 101–110
Integral quantitative assessment, **5**, 10–14
Intertidal zone, **3**, 15–29
Isochaetides resorptus sp. n., **7**, 104–107

Ketone bodies formation in fish, **5**, 101–106
Kremenchug Reservoir, **4**, 92–102; **5**, 36–43

Lake Baikal, **2**, 71–89; **4**, 70–76; **6**, 93–106
Lake Khanka, **4**, 20–24
Laptev Sea, **6**, 79–83
Leptolida, **6**, 84–92
Lesquereusiidae, **7**, 19–32
Lie-in-wait behavior, **4**, 25–33
Long-lived radionuclide accumulation, **5**, 107–110
Lower Danube, **1**, 92–101

Macerrima complex, **7**, 78–84
Macrobenthos zonal distribution, **4**, 12–19

Macrophyte vegetation alteration, **1**, 11–28
Mantle complex evolution, **7**, 51–58
Marine environment, **2**, 49–58
Marine nematode, **4**, 20–24
Marine trematodes, **4**, 25–33
Meiobenthos migration, **1**, 45–54
Metabolism of organic compounds, **2**, 40–48
Microelement composition, **6**, 1–8
Microsporidia, **6**, 61–65
Microsporidians, **6**, 61–65
Mnemiopsis ctenophores, **3**, 9–14
Mollusca, **7**, 8–18
Mollusks, **5**, 107–110
Morphological features, **7**, 78–84
Morphology, **2**, 94–108
Mud-dwelling mussels, **1**, 66–75
Muddy biotopes, **5**, 30–35
Mussel mariculture, **2**, 49–58
Mytilus galloprovincialis, **5**, 30–35

Neanthes succinea, **2**, 23–32
Nematoda, **4**, 20–24; **6**, 76–78
Nemertea, **6**, 22–34
Neuston algae, **1**, 1–10
New genus, **4**, 65–69
New species, **4**, 20–24; 65–69; **7**, 19–32; 78–84; 99–103
Northern Eurasia, **7**, 8–18
Northwest Pacific, **3**, 90–96
Northwestern Black Sea, **2**, 23–32
Nudibranch family, **7**, 59–67
Nudibranchia, **7**, 51–58
Nudibranchs, **7**, 51–58

Octoporia species, **3**, 1–8
Oligochaeta, **4**, 70–76; **7**, 104–107
Opisthobranchia, **7**, 59–67
Organic-pollution indicators, **7**, 4–7
Origin of free-living Turbellaria, **2**, 71–89
Oxystominidae, **4**, 20–24

Pandalidae, **6**, 79–83
Pandalus borealis larvae, **6**, 79–83
Parasitic hydroids, **7**, 1–3
Paraugaptilus, **7**, 99–103
Pelagic waters, **2**, 23–32
Pelmatohydra oligactis, **7**, 85–98
Pentactula larvae, **6**, 35–44
Periphytic biocenoses, **5**, 44–50
Periphyton algae, **1**, 84–91
Periphyton communities, **2**, 13–22
Periwinkles, **5**, 80–90
Phosphorus regime, **1**, 92–101
Phyllophorus field, **2**, 23–32
Phylogenetic relationships, **6**, 61–65
Physiological response, **6**, 66–75
Phytoplankton abundance, **1**, 101–110
Phytoplankton community, **5**, 10–14
Phytoplankton community structure, **5**, 10–14
Phytoplankton effect on pH, **5**, 15–29
Pisidioidea, **7**, 8–18
Plankton, **4**, 103–109
Planktonic infusoria, **7**, 4–7
Polychaeta, **3**, 90–96; 97–113; **4**, 34–50
Polychaete, **2**, 23–32
Polynoidae, **3**, 97–113
Polyurethane foam, **1**, 36–44
Prayidae, **7**, 33–50
Prayinae, **7**, 33–50
Pripyat River, **7**, 4–7
Problems of hydroecology, **3**, 30–34
Progeny, **3**, 46–53
Proteolytic activity stimulation, **2**, 59–66
Protozoa, **1**, 36–44; **6**, 61–65
Psetta maeotica, **6**, 76–86
Pycnogonida, **7**, 78–84

Radiological conditions, **3**, 54–69
Rare species, **7**, 19–32
Reproduction, **3**, 9–14
Reproduction of *Abramis brama*, **3**, 46–53
Reproductive behavior, **1**, 29–35
Reproductive period, **1**, 66–75
Reservoirs, **6**, 93–106
Review, **6**, 22–34; 45–60
Revision, **7**, 59–67
Revolving disk, **5**, 44–50
Rhizopoda, **7**, 19–32
Rhyacodrilus stephensoni, **4**, 70–76
Rising water level, **3**, 87–89
Rosacea, **7**, 33–50

Rosette diagrams, **3**, 70–75
Rotifer participation, **2**, 40–48

Sanitation, **2**, 49–58
Saprophytic microflora, **5**, 51–65
Saratov Reservoir fishes, **6**, 18–21; 107–110
Sarcodina, **4**, 51–64; **6**, 45–60
Scophthalmidae, **6**, 76–86
Sea of Japan, **2**, 67–70
Seasonal abundance, **1**, 36–44
Seasonal changes, **6**, 1–8
Seasonal dynamics, **5**, 30–35
Self-purification analysis, **5**, 1–9
Self-purification assessment, **5**, 1–9
Self-purification processes, **5**, 1–9
Septibranchia, **3**, 1–8
Sevastopol vicinity, **6**, 76–86
Shelled amoebas, **2**, 90–93; **7**, 19–32
Silver nitrate impregnation, **2**, 90–93
Simocephalus, **7**, 68–77
Siphonophora, **7**, 33–50
Sosane holthei sp. n., **3**, 90–96
Southern coast of Vietnam, **3**, 97–113
Southwest Indian Ocean, **7**, 99–103
Spawning, **5**, 30–35
Species composition, **6**, 45–60
Species review, **7**, 8–18
Spectral technique, **2**, 6–12
Spionidae, **4**, 34–50
Sponge distribution, **3**, 15–29
Sponge population structure, **3**, 15–29
Squid, **5**, 91–100
Stichopodidae, **3**, 97–113
Study method, **2**, 90–93
Subgenus *Euglesa*, **7**, 8–18
Survival rate, **3**, 46–53

Testacealobosia, **7**, 19–32
Tissue chemistry, **1**, 66–75
Tissue lipid composition, **5**, 91–100
Tobrilidae, **6**, 76–78
Tobrilus wesenbergi, **6**, 76–78
Trace elements, **2**, 6–12
Trematoda, **6**, 18–21; 107–110
Trematode, **6**, 107–110
Trematode size structure variation, **6**, 18–21
Tropical Atlantic, **5**, 91–100
Tubificidae, **4**, 70–76; **7**, 104–107
Tubularia larynx, **6**, 84–92
Tubulariidae, **6**, 84–92
Turbot, **6**, 76–86
Tyulka, **5**, 36–43
Tyul'ka diet, **4**, 92–102

Ukraine, **4**, 103–109

Vil'kitskiy Strait, **6**, 79–83
Volga River, **3**, 54–69

Warm-water rearing, **6**, 1–8
Water treatment, **5**, 44–50
Water-quality determination, **5**, 66–79
Water-quality evaluation, **3**, 70–75
Western Bug River, **3**, 70–75
White Sea, **4**, 12–19; 51–64; **5**, 80–90

Yenisey reservoirs, **4**, 77–91

Zernova, **2**, 23–32
Zooneuston, **4**, 77–91